



CALCIUM

CALCIUM (Ca) IN SOILS

Calcium is an important plant nutrient. It also plays an important role in determining soil physical and chemical parameters, i.e. structure and pH.

Calcium ions cause soil colloids (clay platelets) to bond or aggregate together, forming crumbs or peds. Soils dominated by calcium are friable and well-structured, have good internal drainage, and are easy to cultivate. They are often described as self-mulching.

In contrast sodium (Na), and to a lesser extent magnesium (Mg), cause clay platelets to disperse. Soils dominated by Na and/or Mg have low infiltration rates, crust after rain, are puggy when wet, set hard on drying and are difficult to cultivate.

Exchangeable calcium levels and soil acidity are usually closely related. Calcium is most available in the pH range 7.0 to 8.5. Under low pH or acid soil conditions, exchangeable calcium levels in the soil are usually low, and the solubility of manganese (Mn) and aluminium (Al) increase and may become toxic.

CALCIUM IN PLANTS

Plants take up calcium in the ionic form (Ca^{2+}). Uptake is not as efficient as for other plant nutrients. It occurs just behind the root tip, in contrast with potassium where uptake occurs along most of the length of the root. Consequently, anything that affects new root growth may prevent calcium uptake and induce a deficiency. This includes adverse weather conditions such as drought, low temperatures, high humidity, poor soil aeration and water logging. For example, "blossom-end" rot in tomatoes, which is attributed to inadequate calcium, can be induced by a period of moisture stress, even though the soil may have adequate calcium levels.

Competition also occurs with other positively charged cations, e.g. ammonium (NH_4^+), potassium (K^+) and magnesium (Mg^{2+}) for root uptake.

Within plants, calcium is not mobile. Once calcium is deposited in leaves, it cannot be remobilised from them to the growing tips. Calcium is required for cell elongation and cell division. Adequate calcium helps delay leaf senescence and slows down or prevents leaf and fruit fall (abscission).

DEFICIENCY SYMPTOMS

Since plants are unable to utilise calcium from old leaves, deficiency normally occurs first in the growing points and youngest leaves. Roots are usually affected before the tops, with both roots and tops exhibiting die back of the growing point. Where Ca deficiency is moderate to acute, root growth is markedly impaired, and plants become susceptible to root-rot infection. Deficiency symptoms also occur in fruit or storage tissues. Some specific symptoms of calcium deficiency are:

- "Bitter pit" in apples - small brown necrotic spots (2 – 3 mm in depth and diameter) over the surface of the fruit.

- “Blossom-end rot” in tomatoes - breakdown at the flower end of the fruit, with depressed blackened patches which may be up to 5 cm wide.
- “Black heart” in celery - deformed and chlorotic (yellow) growth. At a more advanced stage the leaf margins become necrotic, i.e. dead patches will be evident at the edges.
- “Pops” in peanuts - empty shells or small kernels in the pods.

CALCIUM FERTILISERS AND SOIL AMENDMENTS

Several factors must be considered in assessing the need to apply calcium and the form and rate at which it is best applied. These include whether the product is being applied as a soil ameliorant or as a calcium fertiliser, the solubility and price of the calcium compound, and the susceptibility of the crop to calcium deficiency.

Soil ameliorants, such as lime and gypsum, are low priced materials that are supplied and spread in bulk at high rates. They are used on:

Acid (low pH) Soils

In very acid soils calcium will need to be applied in a form which will correct acidity. **Lime** (calcium carbonate) or some other liming material needs to be used. Typical application rates for lime are in the range of 2.5 to 7.5 t/ha.

Lime is insoluble and takes time to react in the soil. In annual crops it should be applied several months ahead of planting and be incorporated into the soil, e.g. at the start of the fallow period. Timing is less critical in tree crops where it is normally applied in winter, in advance of the spring flush and the main growing season.

Being insoluble the effectiveness of lime is very dependent on its hardness and particle size. Hard limestone that is coarser than 250 microns (0.25 mm) has little value in raising soil pH, at least in the short term. Soft limes are more reactive.

Poorly structured (Sodic) soils

Unless the soil is also acidic ($\text{pH}_w < 6.5$), in which case lime can be used, the soil ameliorant that should be used to improve the structure of sodic soils is **Gypsum** (calcium sulphate). Application rates are typically in the range of 5 – 10 t/ha.

While only sparingly soluble, gypsum is much more soluble than lime allowing it to be used on all soil types, irrespective of the pH. Lime is unsuitable for use on neutral to alkaline soils as it is insoluble. Lime will not dissolve and react in the soil unless the soil is acid.

Like lime, gypsum needs to be applied well ahead of planting and to be thoroughly incorporated into the soil. This allows time for calcium to displace sodium from the surface of clay platelets in the soil, and to be leached deeper into the soil. Gypsum has little or no effect on soil pH. It will not correct soil acidity.

Calcium often has to be applied in fertiliser programs during the growing season in fruit and vegetable crops that are susceptible to calcium deficiency, even on soils that are well endowed with calcium or that have been recently limed, as a preventative measure against deficiency.

Calcium Nitrate, while more costly than other calcium products, is recommended in such circumstances. Calcium nitrate has high solubility in water, so is the product of choice where a quick

response to calcium is required. It can be applied in fertigation programs and/or as foliar sprays. Routine applications are required every one to two weeks during critical growth stages, e.g. periods of rapid growth and fruit filling. Deficiency should not be allowed to develop before commencing application.

Calcium nitrate may also be used as a non-acidifying nitrogen fertiliser through drip and trickle irrigation systems and under tree sprinklers.

SuPerfect® (Single Superphosphate) also contains 19% Calcium. SuPerfect is used to supply phosphorus (P). Its use can also contribute to meeting the calcium requirement of crops and pastures where the soil is low in calcium.

SuPerfect is comparatively high in cadmium (Cd) compared to the high analysis phosphorus fertilisers marketed by Incitec Pivot Fertilisers, e.g. DAP. Planting fertilisers that are low in Cd should be used in vegetables. It is recommended that SuPerfect not be used as the sole source of phosphorus in NPK fertiliser blends for vegetables.

SAFETY DIRECTIONS

Refer to the Safety Data Sheet (SDS) for more detailed safety advice. Before use, read the Product Label and the SDS. Use safe work practices and avoid contact with the eyes and skin. Avoid ingestion and inhaling dust. Protective clothing, eyewear and dust masks should always be used when dealing with this product. Observe good personal hygiene, including washing hands after use. Avoid loss of fertiliser to waterways.

WARNING

This document contains information of a general nature. Before using fertiliser seek independent agronomic advice. Fertiliser programs may need to be varied depending on the plants being grown, climatic and soil conditions, application methods, irrigation, agricultural and livestock management practices, the soil's fertility, and cultural practices. ('Unforeseen Elements')

Fertiliser may burn and/or damage crop roots or foliage. Foliar burn to the leaves, fruit or other plant parts is most likely to occur when fertilisers are foliar applied at high concentrations and/or on a regular basis, different products are mixed and sprayed together at cumulatively high rates, the water is of poor quality, or the spray is applied under hot dry conditions, e.g. in the heat of the day.

Fertiliser and supplements may affect animal health. Seek independent advice before using any supplements in livestock rations.

DISCLAIMER

As Unforeseen Elements are beyond the control of Incitec Pivot Limited, in no event Incitec Pivot Limited and its related bodies corporate be liable or accept any responsibility whatsoever for any direct, indirect, punitive, incidental, special or consequential damages (including but not limited to loss of revenue, crops and livestock), in respect of the illness, injury or death of a person, damage to property (including of a third party), or any other loss whatsoever arising out of or connected with the use or misuse of this fertiliser, or its transport, storage, handling or application. Where Incitec Pivot Limited and its related bodies corporate's liability cannot be lawfully excused, it and its related bodies corporate's liability shall be limited to the replacement of, or cost of the fertiliser supplied. The buyer accepts and uses this product subject to these conditions.



COPYRIGHT

Copying or reproduction in whole, or in part, by any means, or transmission, or translation into a machine language without the written permission of Incitec Pivot Limited, is strictly prohibited.