



# COMPLETE BALANCED NUTRITION IN A SINGLE GRANULE.

**GRANULOCK<sup>®</sup>**  **BLUE**

[incitecpivotfertilisers.com.au](http://incitecpivotfertilisers.com.au)

UNIFORM  
NUTRIENT  
DELIVERY  
TO HELP  
**MAXIMISE  
CROP  
PRODUCTION  
AND QUALITY.**

**GRANULOCK** 

## WHY CHOOSE GRANULOCK BLUE?

### BALANCED NUTRITION IN EVERY GRANULE

Each granule contains all the nutrients, improving the uniformity of nutrient distribution in crops and reducing issues with nutrient segregation during transport or application.

### OPTIMAL PLANT ESTABLISHMENT

The nitrate nitrogen and water-soluble form of the key nutrients included in Granulock Blue allow quick plant uptake for vigorous plant establishment.

### EVENLY-SIZED GRANULES FOR EVEN RESULTS

The granules are uniformly sized for even spreader application, promoting even crop growth and produce quality.

### AMMONIUM STABILISER TREATABLE

Nutrient efficiency can be improved and leaching, runoff and denitrification pathways reduced.

### RAPID RELEASE WITH MINIMAL MOISTURE

For improved nutrient access the water-soluble nutrients are rapidly released into the soil following night dew or as little as 3mm of rain.

### REDUCED N LOSSES THROUGH VOLATILISATION

The favourable pH of the actual fertiliser granule coupled with having nitrogen present in the nitrate and ammonium forms, reduces the potential for volatilisation losses when Granulock Blue is surface-applied compared with urea blends.

### EASY APPLICATION AND STORAGE

The high-quality granules are resin-treated for free-flowing, precise application and ease of storage.

Granulock Blue is a versatile compound NPKS fertiliser. It provides uniform delivery of targeted, balanced nutrition to maximise yield potential and quality in tree, vine and vegetable crops, as well as sugar cane. Granulock Blue also promotes vigour in floriculture ornamental plants and turf.

## ANALYSIS

Nitrogen	Ammonium Form	7.1% w/w
Nitrogen	Nitrate Form	4.9% w/w
Nitrogen	Total	12.0% w/w
Phosphorus	Water Soluble	2.6% w/w
Phosphorus	Citrate Soluble	2.6% w/w
Phosphorus	Total	5.2% w/w
Potassium	as Sulfate	14.1% w/w
Sulfur	as Sulfate	8.0% w/w
Calcium	as Sulfate / Phosphate	3.6% w/w
Magnesium	as Magnesium Sulfate	1.2% w/w
Boron	as Sodium Borate	0.02% w/w

## IMPURITIES

Cadmium	Maximum	1mg/kg
Cadmium	Maximum	10mg/kg P
Lead	Maximum	10mg/kg P
Mercury	Maximum	0.2mg/kg

## GETTING THE BEST OUT OF GRANULOCK BLUE

### GRANULOCK BLUE IS RECOMMENDED:

- As a planting fertiliser for vegetables where soil phosphorus levels are moderate to high, or pre-plant phosphorus has been applied.
- As a base or side dressing for intensively grown crops.
- To fertilise chloride-sensitive crops.
- In tree crops, where a broadcast NPKS application is required.
- To fertilise crops grown in salt-affected soils or on soils with a moderate supply of phosphorus.

**GRANULOCK** 





## APPLICATION RECOMMENDED GUIDELINES BY CROP



Vegetables					
Crop	Rate of applications			No. of appl.	Timing
	kg/ha	g/m <sup>2</sup>	g/m of row		
Tomatoes	500 kg/ha	50 g/m <sup>2</sup>	50 (1.5m row)	1	Preplant or planting
Potatoes	500 kg/ha	50 g/m <sup>2</sup>	40 (0.75m row)	1	Preplant or planting
Cucurbits (Pumpkins, Watermelons, Cucumbers, Zucchini)	350 kg/ha	35 g/m <sup>2</sup>	70 (2m row)	1	Preplant or planting
Beans	350 kg/ha	35 g/m <sup>2</sup>	25 (0.75m row)	1	Preplant or planting
Brassicas (Broccoli, Cauliflower, Cabbage)	400-600 kg/ha	45 g/m <sup>2</sup>	30 (0.75m row)	1	Preplant or planting
Leafy crops (Spinach, Rocket, Sorrel)	450 kg/ha	45 g/m <sup>2</sup>	30 (0.75m row)	1	Preplant or planting
Carrots	300 kg/ha	30 g/m <sup>2</sup>	22 (0.75m row)	1	Preplant or planting
Beetroot, Turnips	500 kg/ha	50 g/m <sup>2</sup>	35 (0.75m row)	1	Preplant or planting

Fruit Crops			
Crop	Rate of applications	No. of appl.	Timing
Stone fruit (High chill) Bearing trees (1,000 trees/ha)	400-650 g/tree	1	Early spring
Stone fruit (Low chill) Bearing trees (500 trees/ha)	500 g/tree	1	July/August (bud break)
	110 g/tree	1	December (post harvest)
	500 g/tree	1	February (hardening of summer flush)
Apples Bearing trees (1,000 trees/ha)	350 g/tree/application	2	Bud burst and late spring
Grapes	250 - 400 g/vine	1	Late winter before bud burst
Turf/Lawns	250 kg/ha or 25 g/m <sup>2</sup>	2	September and January
Palms Large, established and bearing	300 g/tree/application	4	October, December, February, April
Flowers (Annuals)	2 level tablespoons/metre of row	1	Before planting
Flowers (Roses)	4 level tablespoons/m <sup>2</sup> of bed	1	At planting
	2 level tablespoons/m <sup>2</sup> of bed	1	August
	1 level tablespoon/m <sup>2</sup> of bed	1	When buds are forming
Flowers (Camellias, Azaleas)	2 level tablespoons/m <sup>2</sup> /application	2	Late August, February
Flowers (Bulbs, Corms)	4 level tablespoons/m <sup>2</sup>	1	3 weeks before planting

Sugar Cane			
Sugar Cane	375 - 450 kg/ha	1	At planting

Tree and Vine Crops			
Crop	Rate of application	No. of applications	Timing
Citrus	400 g/tree x age in years (up to 3.2kg/tree at 8 years and older)	1	July
Paw Paw	30 g/tree/month	4-5	From planting to flowering
	60 g/tree/month	5-6	From flowering to first harvest
	175 g/tree	2	In the first month following harvest and again six months later for the second and subsequent years
Macadamia	100 g/tree/application	5 in year 1	February, April, June, October, December
	150 g/tree/application	5 in year 2	February, April, June, October, December
	350 g/tree/application	5 in year 3	February, April, June, October, December
	425 g/tree/application	5 in year 4	February, April, June, October, December
	500 g/tree/application	5 in year 5	February, April, June, October, December
	1040 g/tree/application	5 in year 6	March, June, August
	1300 g/tree/application	5 in year 7	March, June, August
Up to a maximum of 6.5 kg/tree/year at 12 years and older			
Avocado Trees 3 years and older For a canopy diameter of 2 metres	140 g/tree/application	2	After fruit drops and 6 weeks later
For a canopy diameter of 4 metres	75 g/tree	1	Early April
	625 g/tree/application	2	After fruit drops and 6 weeks later
For a canopy diameter of 6 metres	285 g/tree	1	Early April
	1250 g/tree/application	2	After fruit drops and 6 weeks later
For a canopy diameter of 8 metres	625 g/tree	1	Early April
	2250 g/tree/application	2	After fruit drops and 6 weeks later
	1070 g/tree	1	Early April
Bananas Plant crop - planting	400 g/plant	1	Preplant incorporated
	200 g/plant	1	In drill at planting (avoid direct contact with planting material)
Plant crop - sidedress	100-250 g/plant/application	8-10	Every 4-6 weeks post planting
Ratoon crop	200 g/plant/application	4	Strategic applications in conjunction with regular fertiliser program



## FIND OUT EXACTLY WHAT NUTRITION YOUR CROP NEEDS

Only with regular and routine use of strategic nutrient analysis (including soil, plant tissue and water analysis, if irrigating) can you tell just what you're missing, and how much you need to apply for optimum crop efficiency and yield.

Let Nutrient Advantage® lend a hand. For over 60 years, our Nutrient Advantage service has been helping farmers gain a productivity edge through reliable soil and plant tissue testing, and expert, localised nutrient advice.

### WANT THE ADVANTAGE?

Visit [nutrientadvantage.com.au](http://nutrientadvantage.com.au) or call **1800 803 453** and ask about our soil and plant tissue testing services.

Other fertilisers may also be needed to support healthy crop growth. Nutrient Advantage® soil testing will assist in monitoring soil nutrient levels and crop requirements. Note: Planting fertilisers should be banded 5 cm to the side and 5 cm below the seed or transplants, and not placed in direct contact with the planting material.

While the fertiliser rates and programs detailed in the table above are a practical guide to good agronomic practice under most situations, they offer indicative rates only and are not complete fertiliser programs. Local soil, climatic and other conditions should be taken into account. Higher or lower rates may be required or different fertilisers altogether. Talk to your local Icitec Pivot Fertilisers Distributor about soil testing and arranging a specific fertiliser recommendation that best suits your needs. This table covers use recommendations for some agricultural segments and is not necessarily a complete guide.



# GRANULOCK<sup>®</sup> BLUE

- ✓ Balanced nutrition in every granule
- ✓ Fast uptake and excellent availability for optimal plant establishment and growth
- ✓ Evenly sized granules for uniform application and even growth
- ✓ Chloride-free potassium for use on salt-affected soils and chloride-sensitive crops

**For more information about Granulock Blue contact your  
Incitec Pivot Fertilisers Distributor or go to [incitecpivotfertilisers.com.au](http://incitecpivotfertilisers.com.au)**



®Fertcare is a registered trademark of Australian Fertiliser Services Association, Inc.  
®Nutrient Advantage is a registered trademark of Incitec Pivot Limited. Incitec Pivot Fertilisers is a business of Incitec Pivot Limited, ABN 42 004 080 264.  
This is a guide only, which we hope you find useful as a general tool. While Incitec Pivot Fertilisers has taken all reasonable care in the preparation of this guide, it should not be relied on as a substitute for tailored professional advice and Incitec Pivot Fertilisers accepts no liability in connection with this guide.

